

SA53A-09: Results from a new approach to meso-scale driving of the I-T system: HIME

Dogacan Su Ozturk¹

Xing Meng¹, Olga Verkhoglyadova¹, Roger Varney², Ashton Reimer², Josh Semeter³


- 1: Jet Propulsion Laboratory, California Institute of Technology:
- 2: Center for Geospace Studies, SRI International, Menlo Park
- 3: Center for Space Physics, Boston University



Role of Meso-scale Electric Fields in I-T Energy Budget

$$Q_{JH} = \sigma_P [\boldsymbol{E} + u_n \times \boldsymbol{B}]^2$$

- Traditionally, conductivity and electric fields are estimated through empirical models.
- Current models can not resolve the meso-scale structures but significant work is ongoing*.




^{*[}Codrescu et al. 2008; Deng et al. 2009; Matsuo & Richmond 2008; Zhu et al. 2018].

Role of Meso-scale Electric Fields in I-T Energy Budget

$$Q_{JH} = \sigma_P [\mathbf{E} + u_n \times \mathbf{B}]^2$$

- Traditionally, conductivity and electric fields are estimated through empirical models.
- Current models can not resolve the meso-scale structures but significant work is ongoing*.




^{*[}Codrescu et al. 2008; Deng et al. 2009; Matsuo & Richmond 2008; Zhu et al. 2018].

Role of Meso-scale Electric Fields in I-T Energy Budget


$$Q_{JH} = \sigma_P [\boldsymbol{E} + u_n \times \boldsymbol{B}]^2$$

 Traditionally, conductivity and electric fields are estimated



Meso-scale electric fields (<15 minutes, 150-500 km) introduce as much energy as the large-scale structures.

resolve the meso-scale structures but significant work is ongoing*.




^{*[}Codrescu et al. 2008; Deng et al. 2009; Matsuo & Richmond 2008; Zhu et al. 2018].


High-latitude Input for Meso-scale Electrodynamics: HIME

Nicolls et al. (2014)
 technique estimates Fregion electric fields on a
 2-D grid from the PFISR
 V_{I OS} measurements.

 These estimates can be merged with a global empirical potential model to drive global I-T models and investigate the role of meso-scale electric fields.



Running a global I-T model with HIME



<u>Modelling:</u>


- PFISR aiding the ISINGLASS experiment with 15 beams operating [Clayton et al., 2019a, 2019b, JGR.]
- The electric field estimates:
 - Temporal resolution [66 seconds]
 - Spatial resolution [0.05° in latitude and 0.3° in longitude, ~10 km]
 - Down-sampled to 0.75°x0.75°: 35x80 km

Validation:

- PFISR plasma measurements along Beams
- Three SDI (Scanning Doppler Imagers) measurements for neutral wind

6 jpl.nasa.gov

Running a global I-T model with HIME




HIME is an add-on framework and does not require major source code modification.


Global Ionosphere -Thermosphere Model (GITM)

- 3D, altitude based non-uniform grid, assumes non-hydrostatic solution
- High-latitude input: Electric potential and particle precipitation
- Output: Plasma and neutral density, temperature, ion and electron velocity, neutral winds


Ridley, Deng and Toth, JASTP, 2006




Electric field estimates from PFISR successfully incorporated.




Estimates of velocity improved for all three components. jpl.nasa.gov




Electric field estimates from PFISR successfully incorporated.




Estimates of velocity improved for all three components. jpl.nasa.gov




Electric field estimates from PFISR successfully incorporated.



Estimates of velocity improved for all three components. jpl.nasa.gov





Electric field estimates from PFISR successfully incorporated.



Estimates of velocity improved for all three components. jpi.nasa.gov

Modeled ion and electron temperature response






Modeled ion temperature <u>captures</u>
<u>variability</u> better when meso-scale electric
fields are used.

Modeled electron temperature <u>captures</u> the <u>amplitude</u> but does <u>not reproduce the short-lived peaks</u>.


Modeled ion and electron temperature response




Modeled ion temperature <u>captures</u>
<u>variability</u> better when meso-scale electric
fields are used.

Modeled electron temperature <u>captures</u> the <u>amplitude</u> but does <u>not reproduce the short-lived peaks</u>.


Modeled electron density response






Poker Flat DASC, collocated with PFISR, shows bright dynamic aurora over the same region.


Modeled electron density response






Increase in simulated electron density likely due decreased westward transport mechanism.

Energy transfer rate with meso-scale electric fields





The deposited energy increases locally in HIME-driven simulation compared to Weimer-driven simulation above 110 km.

Lotko and Zhang, 2019

Effects of Meso-scale electric fields in the I-T system

The HIME driven simulations showed that meso-scale electric fields can lead to:

- Variations in the electron density due to transport.
- Collisional and frictional heating of the plasma and neutrals.
- Increased volumetric heating along the vertical profiles.

Ongoing and future work on HIME:

- A self-consistent treatment of particle precipitation is necessary to fully understand the effects of meso-scale structures.
- Further case studies are necessary to assess the performance of HIME.

Thank you.

Acknowledgements

- This work is funded by the NASA ROSES 2016 Heliophysics LWS Science (NRA NNH16ZDA001N) Program.
- GITM is developed and supported by Prof. Aaron Ridley at University of Michigan.
- Simulations were done on NASA High-End Computing Program through the NASA Advanced Supercomputing Division at Ames Research Center, and Stampede Supercomputer at Texas Advanced Computing Center at University of Texas at Austin.
- We acknowledge use of NASA/GSFC's Space Physics Data Facility's OMNIWeb (or CDAWeb or ftp) service, and OMNI data.
- We acknowledge NASA contract NAS5-02099 and V. Angelopoulos for use of data from the THEMIS Mission. Specifically: C. W. Carlson and J. P. McFadden for use of ESA data; K. H. Glassmeier, U. Auster and W. Baumjohann for the use of FGM data provided under the lead of the Technical University of Braunschweig and with financial support through the German Ministry for Economy and Technology and the German Center for Aviation and Space (DLR) under contract 50 OC 0302.
- Simulation results have been provided by the Community Coordinated Modeling Center at Goddard Space Flight Center through their public Runs on Request system (http://ccmc.gsfc.nasa.gov). The Weimer Model was developed by Daniel R. Weimer at Virginia Tech. The Ovation Prime Model was developed by Patrick Newell at JHU/APL.
- Jeffrey Nee helped access to visual material presented in this seminar.



jpl.nasa.gov